

FINITE JET DETERMINATION OF CONSTANTLY DEGENERATE CR EMBEDDINGS

PETER EBENFELT AND BERNHARD LAMEL

ABSTRACT. We prove finite jet determination results for smooth CR embeddings which are of constant degeneracy, using the method of complete systems. As an application, we derive a reflection principle for mappings between a Levi-nondegenerate hypersurface in $\mathbb{C}^N$ and a Levi-nondegenerate hypersurface in $\mathbb{C}^{N+1}$. We also give an independent proof of the reflection principle for mappings between strictly pseudoconvex hypersurfaces in any codimension due to Forstneric [6].

1. INTRODUCTION

In this paper, we shall derive a complete system of differential equations for CR mappings between a smooth hypersurface in $\mathbb{C}^N$ and a smooth hypersurface in $\mathbb{C}^{\hat{N}}$ (see e.g. Theorem 8). This system of differential equations will then be used to prove finite determination results (Theorems 1, 2, and 7) for such mappings. For historical information and background on the finite determination problem, we refer the reader to the introduction of [5] or the survey article [1]. As an application of our main results, we also prove a new reflection principle for smooth mappings between real-analytic Levi nondegenerate hypersurfaces (Corollary 5).

To formulate our results precisely, we shall need to introduce some notation. Let M be a real hypersurface in $\mathbb{C}^N$, $N \geq 2$, and let us denote by $\mathcal{V} \subset \mathbb{C}TM$ its CR bundle; the reader is referred e.g. to [2] for basic notions and facts about CR manifolds. Recall that a mapping $f: M \rightarrow \mathbb{C}^k$ is called CR if $f_*(\mathcal{V}) \subset T^{0,1}\mathbb{C}^k$, where f_* denotes the induced mapping of the tangent spaces (the push forward). This is equivalent to saying that, near every point $p \in M$, there is a neighborhood $U \subset M$ of p in which

$$(1) \quad L_{\bar{a}}f_j = 0, \quad a = 1, \dots, n, \quad j = 1, \dots, k,$$

where $n = N - 1$ and $L_{\bar{1}}, \dots, L_{\bar{n}}$ is a basis for the CR vector fields in U . We use here the convention that e.g. $L_{\bar{a}} = \overline{L_a}$ so that the $L_1, \dots, L_n$ form a basis for the vector fields valued in $\overline{\mathcal{V}}$ near p on M .

Suppose that $\hat{M} \subset \mathbb{C}^{\hat{N}}$ is a real hypersurface and $f: M \rightarrow \mathbb{C}^{\hat{N}}$ is a CR mapping sending M into $\hat{M}$. Let p be a point in M and $\hat{p}$ a local defining function for $\hat{M}$ near $\hat{p} := f(p) \in \hat{M}$. The mapping f is called *transversal* at p if $f_*(\mathbb{C}T_p M)$ is not contained in the subspace (of codimension 1) $\hat{\mathcal{V}}_{\hat{p}} + \overline{\hat{\mathcal{V}}}_{\hat{p}} \subset \mathbb{C}T_{\hat{p}}\hat{M}$. One can show (see section 3.4) that if f is transversal and M is Levi nondegenerate at $p \in M$, then f is in fact a local embedding near p .

Following Lamel [9], we define an increasing sequence of subspaces $E_k(q) \subset \mathbb{C}^{\hat{N}}$, for $q \in M$ near p , associated to the germ of the mapping $f: (M, q) \rightarrow \hat{M}$ as follows. Let $L_{\bar{1}}, \dots, L_{\bar{n}}$ be a basis for the CR vector fields on M near p as above and define

$$(2) \quad E_k(q) := \text{span}\{L^{\bar{J}}(\hat{p}_{Z'} \circ f)(q) : J \in (\mathbb{Z}_+)^n, |J| \leq k\} \subset \mathbb{C}^{\hat{N}},$$

1991 *Mathematics Subject Classification.* 32H02.

The first author is a Royal Swedish Academy of Sciences Research Fellow supported by a grant from the Knut and Alice Wallenberg Foundation and also by DMS-0100110. The second author was supported by the ANACOGA research network.

where $\hat{\rho}_Z = (\partial\hat{\rho}/\partial Z'_j)_{1 \leq j \leq \hat{N}}$ in some local coordinate system Z' near $\hat{p}$; we use here standard multi-index notation $L^{\bar{J}} := L_1^{\bar{J}_1} \dots L_{\bar{n}}^{\bar{J}_{\bar{n}}}$ and $|J| = J_1 + \dots + J_n$. One can show (cf. [9]) that the increasing sequence of numbers $d_k(q) := \dim E_k(q)$ is independent of the choice of local defining function $\hat{\rho}$ and coordinates Z' , as well as of the choice of basis of the CR vector fields $L_1, \dots, L_{\bar{n}}$. We shall say, again following Lamel (*loc. cit.*) that $f: (M, q) \rightarrow \hat{M}$ is (k_0, s) -degenerate at q if s is the minimum of the decreasing sequence of codimensions of E_k , i.e.

$$s := \min_k (\hat{N} - d_k),$$

and k_0 is the smallest integer such that this minimum is attained. The case where $s = 0$ is actually a nondegenerate situation, and we say that f is k_0 -nondegenerate if it is $(k_0, 0)$ -degenerate. If f is (k_0, s) -degenerate at a point q and the degeneracy s is constant in a neighborhood of q , then we shall say that f is constantly (k_0, s) -degenerate at q . The notion of (k_0, s) -degeneracy will be presented in a more intrinsic form in section 3.2.

The following are the main results in this paper.

Theorem 1. *Let $M \subset \mathbb{C}^N$ and $\hat{M} \subset \mathbb{C}^{N+1}$ be smooth hypersurfaces which are Levi-nondegenerate at $p_0 \in M$ and $\hat{p}_0 \in \hat{M}$, and $f: M \rightarrow \hat{M}$ a smooth CR mapping with $f(p_0) = \hat{p}_0$ which is constantly (k_0, s) -degenerate and transversal at p_0 . Then f is uniquely determined by its $2k_0$ -jet $j_{p_0}^{2k_0} f$ at p_0 in a neighbourhood of p_0 . That is, there exists a neighbourhood U of p_0 in M such that if $g: M \rightarrow \hat{M}$ is another smooth CR map of constant degeneracy s with $j_{p_0}^{2k_0} f = j_{p_0}^{2k_0} g$, then $f|_U = g|_U$.*

Theorem 2. *Let $M \subset \mathbb{C}^N$ and $\hat{M} \subset \mathbb{C}^{\hat{N}}$ be smooth hypersurfaces, $p_0 \in M$, $\hat{p}_0 \in \hat{M}$, $\hat{M}$ strictly pseudoconvex at $\hat{p}_0$, and $f: M \rightarrow \hat{M}$ a smooth CR mapping with $f(p_0) = \hat{p}_0$ which is constantly (k_0, s) -degenerate at p_0 . Then f is uniquely determined by its $2k_0$ -jet $j_{p_0}^{2k_0} f$ at p_0 in a neighbourhood of p_0 . That is, there exists a neighbourhood U of p_0 in M such that if $g: M \rightarrow \hat{M}$ is another smooth CR map of constant degeneracy s with $j_{p_0}^{2k_0} f = j_{p_0}^{2k_0} g$, then $f|_U = g|_U$.*

Let us note here that in the setting of Theorem 2, the mapping f is automatically transversal (see e.g. the paper [6] by Forstneric). The reader should also notice that in Theorem 2, where the hypersurfaces are required to be strictly pseudoconvex, there is no restriction on the codimension $\hat{N} - N$ of the mapping f whereas in Theorem 1, where no pseudoconvexity is assumed, the codimension is restricted to be one. The reason for this is easily seen by considering the case where $M \subset \mathbb{C}^N$ is given by the equation

$$\Im z_1 = \sum_{k=2}^N |z_k|^2$$

and $\hat{M} \subset \mathbb{C}^{N+2}$ by

$$\Im z_1 = \sum_{k=2}^N |z_k|^2 + |z_{N+1}|^2 - |z_{N+2}|^2.$$

The mapping

$$(3) \quad (z_1, \dots, z_N) \mapsto (z_1, \dots, z_N, f(z), -f(z)),$$

where f is an arbitrary holomorphic function with $f(0) = 0$, sends $(M, 0)$ into $(\hat{M}, 0)$. It is not difficult to see that if k_0 is the smallest integer ≥ 2 for which there is a multi-index $\alpha \in \mathbb{Z}_+^{N-1}$ with $\partial_z^\alpha f(0) \neq 0$, where $z' = (z_2, \dots, z_N)$, then the mapping (3) is constantly $(k_0, 1)$ -degenerate at 0. Clearly, you can find two different such functions f_1, f_2 whose jets at 0 agree up to arbitrary order and, hence, Theorem 1 is false in codimension 2.

We shall see that in fact, given a mapping f as in the preceding theorems, there exists a dense, open subset M_f of M such that for each $p \in M_f$, f is constantly (k_0, s) -degenerate for some s and furthermore, that $k_0 \leq \hat{N} - N + 1 - s \leq \hat{N} - N + 1$ (see Lemma 12). Hence, we have the following corollaries.

Corollary 3. *Let $M \subset \mathbb{C}^N$ and $\hat{M} \subset \mathbb{C}^{\hat{N}+1}$ be smooth hypersurfaces which are Levi-nondegenerate, and $f : M \rightarrow \hat{M}$ and $g : M \rightarrow \hat{M}$ smooth, transversal CR mappings. If for any p_0 in the dense open subset $M_f \cap M_g$ of M we have that $j_{p_0}^{2(\hat{N}-N+1)} f = j_{p_0}^{2(\hat{N}-N+1)} g$, then $f = g$.*

Corollary 4. *Let $M \subset \mathbb{C}^N$ and $\hat{M} \subset \mathbb{C}^{\hat{N}}$ be smooth strictly pseudoconvex hypersurfaces and $f : M \rightarrow \hat{M}$ and $g : M \rightarrow \hat{M}$ smooth CR mappings. If for any p_0 in the dense open subset $M_f \cap M_g$ of M we have that $j_{p_0}^{2(\hat{N}-N+1)} f = j_{p_0}^{2(\hat{N}-N+1)} g$, then $f = g$.*

As an application of Theorem 1, we obtain the following reflection principle.

Corollary 5. *Let $M \subset \mathbb{C}^N$ and $\hat{M} \subset \mathbb{C}^{\hat{N}+1}$ be real-analytic hypersurfaces which are Levi-nondegenerate, and $f : M \rightarrow \hat{M}$ a smooth transversal CR mapping. Then there exists an open dense subset $M_0 \subset M$ such that for any $p \in M_0$, there exists an open neighbourhood V of p in $\mathbb{C}^N$ and a holomorphic map $F : V \rightarrow \mathbb{C}^{\hat{N}}$ such that $F|_{V \cap M} = f$.*

Let us give the (short) proof here. We first point out that given a mapping f as in Corollary 5, there exists a dense, open subset M_0 of M such that for each $p \in M_0$, f is constantly (k_0, s) -degenerate, for some s , in an open neighborhood of p (see Lemma 11); in fact, one may choose M_0 such that f is locally constantly (k_0, s) -degenerate on M_0 with $s \leq \hat{N} - N$ and $k_0 \leq \hat{N} - N + 1 - s \leq \hat{N} - N + 1$ (see Lemma 12). For points $p_0 \in M_0$, we can use a result of the second author ([8], Theorem 6) to conclude that the formal Taylor series of f at p_0 converges (in some neighbourhood V of p_0) to a holomorphic mapping $F : V \rightarrow \mathbb{C}^{\hat{N}}$ with $F(M \cap V) \subset \hat{M}$. By Theorem 1, $F|_M = f$, which proves the corollary.

The same argument (using Lemma 11 and Theorem 2) also gives an independent proof of the following theorem due to Forstneric [6].

Corollary 6. *Let $M \subset \mathbb{C}^N$ and $\hat{M} \subset \mathbb{C}^{\hat{N}+1}$ be real-analytic strictly pseudoconvex hypersurfaces and $f : M \rightarrow \hat{M}$ a smooth CR mapping. Then there exists an open dense subset $M_0 \subset M$ such that for any $p \in M_0$, there exists an open neighbourhood V of p in $\mathbb{C}^N$ and a holomorphic map $F : V \rightarrow \mathbb{C}^{\hat{N}}$ such that $F|_{V \cap M} = f$.*

A special case for which we also derive finite determination results without any conditions on $\hat{N}$ is the case of *nondegenerate* mappings. These have already been investigated by Kim and Zaitsev [10] in the general codimension case. However, our method is a bit different, and we have an easy, independent proof of the following result for nondegenerate mappings between hypersurfaces.

Theorem 7. *Let $M \subset \mathbb{C}^N$ and $\hat{M} \subset \mathbb{C}^{\hat{N}}$ be smooth hypersurfaces, $p_0 \in M$ a point of finite type, and let $f : M \rightarrow \hat{M}$ be a smooth CR mapping which is k_0 -nondegenerate at p_0 . Then f is uniquely determined by its $2k_0$ -jet $j_{p_0}^{2k_0} f$ at p_0 in a neighbourhood of p_0 . That is, there exists a neighbourhood U of p_0 in M such that if $g : M \rightarrow \hat{M}$ is a smooth CR map with $j_{p_0}^{2k_0} f = j_{p_0}^{2k_0} g$, then $f|_U = g|_U$.*

In order to prove Theorems 1, 2, and 7 we will construct complete systems of differential equations for all the cases considered. To formulate the result more precisely, we shall use the following notation. For an open subset $U \subset M$, we shall denote by $J^m(U, \hat{M})$ the space of m -jets of smooth (not necessarily CR) mappings $F : U \rightarrow \hat{M}$ and by $j_p^m F$ the m -jet of such a mapping at $p \in U$ (which we shall think of as the collection of all derivatives of F , in some local coordinate system near p , up to order m). The result needed to prove Theorems 1 and 2 can now be formulated as follows.

Theorem 8. *Let $M \subset \mathbb{C}^N$, $\hat{M} \subset \mathbb{C}^{\hat{N}}$ be smooth hypersurfaces, $p_0 \in M$, $\hat{p}_0 \in \hat{M}$, $f : M \rightarrow \hat{M}$ a transversal smooth CR map with $f(p_0) = \hat{p}_0$ of constant degeneracy (k_0, s) at p_0 , and assume that either $M \subset \mathbb{C}^N$ and $\hat{M} \subset \mathbb{C}^{N+1}$ are Levi nondegenerate at p_0 and $\hat{p}_0$ respectively, or that M and $\hat{M}$ are strictly pseudoconvex at p_0 and $\hat{p}_0$ respectively (with no restrictions on $\hat{N}$). Then there exists a neighbourhood U of p_0 in M and a smooth function $\phi : U \times J^{2k_0+2}(U, \hat{M}) \rightarrow J^{2k_0+2}(U, \hat{M})$ which only depends on M and $\hat{M}$ such that*

$$(4) \quad j_x^{2k_0+3} f = \phi(x, j_x^{2k_0+2} f), \quad x \in U.$$

Furthermore, there exists a neighbourhood V of $j_{p_0}^{2k_0+2} f$ such that if $g : M \rightarrow \hat{M}$ a transversal smooth CR map with $g(p_0) = \hat{p}_0$ of constant degeneracy (k_0, s) at p_0 with $j_{p_0}^{2k_0+2} g \in V$, then (for some neighbourhood U' of p_0)

$$(5) \quad j_x^{2k_0+3} g = \phi(x, j_x^{2k_0+2} g), \quad x \in U'.$$

Theorems 1 and 2 follow from Theorem 8 using standard arguments about uniqueness of solutions of complete systems of differential equations (see e.g. [4]) and the fact that the m -jet of any mapping $f : M \rightarrow \hat{M}$, as in Theorems 1 and 2, at p_0 , for any m (and hence in particular the $2k_0 + 2$ -jet) is determined by its $2k_0$ jet at p_0 ([8], Theorem 8). Thus, to prove Theorems 1 and 2 it suffices to prove Theorem 8. Also, by a remark given earlier, we could drop the assumption of transversality in the strictly pseudoconvex case, since it is satisfied automatically.

The organization of the paper is as follows: In section 2, we review the definition of a CR vector bundle, give definitions of the nondegeneracy conditions introduced above in this context, and introduce some essential notation and tools. Following that, we give the proof of Theorem 7. In the next two sections, we derive some jet reflection identities which we use in section 7 to prove Theorem 8.

2. CR VECTOR BUNDLES

In this section, we define the notion of a CR vector bundle. This notion is not new and has been extensively used in the literature, but we give a thorough definition nonetheless. We also give some examples, and introduce the canonical derivative operator D (which is the same as $\bar{\partial}_b$ in most of the literature). For our purposes, the local formalism of this derivative operator is more important than the global formalism developed alongside it.

2.1. Definition. Let M be a CR manifold. Let us first recall some basic definitions. We are given a subbundle $\mathcal{V}(M) \subset \mathbb{C}TM$ (which we will refer to as the *CR bundle* of M) which satisfies

$$(6) \quad [\mathcal{V}(M), \mathcal{V}(M)] \subset \mathcal{V}(M), \quad \text{and } \mathcal{V}(M) \cap \overline{\mathcal{V}(M)} = \{0\}.$$

In (6), $[\mathcal{V}(M), \mathcal{V}(M)] \subset \mathcal{V}(M)$ means that for all sections X, Y of $\mathcal{V}(M)$, the section $[X, Y]$ of $\mathbb{C}TM$ takes values in $\mathcal{V}(M)$, and 0 stands for the 0 section of $\mathbb{C}TM$. If M is a CR manifold, we also say that M has a CR structure, and sometimes we even refer to $\mathcal{V}(M)$ as a CR structure on M . If $U \subset M$ is open, a section $\bar{L} \in \Gamma(U, \mathcal{V}(M))$ is called a CR vector field over U . A CR function on U is one which is annihilated by all CR vector fields on U . If $M, \hat{M}$ are two CR manifolds and $f : M \rightarrow \hat{M}$ is a map, say of class C^1 , we say that f is CR provided $f_*(\mathcal{V}(M)) \subset \mathcal{V}(\hat{M})$. Let us also recall that if X is a complex manifold, then X has a natural CR structure given by $\mathcal{V}(X) = T^{(0,1)}X$.

We consider a complex vector bundle E over M with projection $\pi : E \rightarrow M$. A vector bundle atlas $\mathcal{A}$ consists of pairs (U_α, ϕ_α) where $\{U_\alpha : \alpha \in A\}$ is an open cover of M , and $\phi_\alpha : \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{C}^k$ is a diffeomorphism such that $\pi|_{\pi^{-1}(U_\alpha)} = \text{proj}_1 \circ \phi_\alpha$, where proj_l denotes the projection on the l th component in a product. The chart change $\phi_\alpha \circ \phi_\beta^{-1} : (U_\alpha \cap U_\beta) \times \mathbb{C}^k \rightarrow (U_\alpha \cap U_\beta) \times \mathbb{C}^k$, is of the form

$\phi_\alpha \circ \phi_\beta^{-1}(x, v) = (x, \phi_{\alpha\beta}(x)v)$, where the *transition functions* $\phi_{\alpha\beta}$ are smooth functions (on $U_\alpha \cap U_\beta$) valued in $GL(\mathbb{C}^k)$ satisfying the usual cocycle conditions (cf. e.g [7]).

We say that $(E, \mathcal{A})$ is a *CR vector bundle* if for $\alpha, \beta \in A$ the transition functions $\phi_{\alpha\beta}$ are CR. We refer to $\mathcal{A}$ as a *CR atlas* for E , and if a CR atlas is fixed, we simply say that E is a CR vector bundle.

A vector bundle chart (U, ϕ) for E is said to be *compatible* with $\mathcal{A}$ if $\phi_\alpha \circ \phi^{-1}(x, v) = (x, \psi_\alpha(x)v)$ where $\psi_\alpha : U \cap U_\alpha \rightarrow GL(\mathbb{C}^k)$ is CR for all $\alpha \in A$; we say that (U, ϕ) is a *CR chart* (for $(E, \mathcal{A})$), and that U is a *trivializing neighbourhood*. If $\mathcal{A}$ has the property that every CR chart (U, ϕ) for $(E, \mathcal{A})$ belongs to $\mathcal{A}$, we say that $\mathcal{A}$ is a *maximal CR atlas* (for E). By including all compatible charts we can associate to $\mathcal{A}$ a unique maximal atlas $\mathcal{A}_{max}$ with the property that $\mathcal{A} \subset \mathcal{A}_{max}$. Two CR vector bundles $(E, \mathcal{A})$ and $(E, \mathcal{A}')$ are compatible if $\mathcal{A}_{max} = \mathcal{A}'_{max}$. Equivalently, the CR vector bundles $(E, \mathcal{A})$ and $(E, \mathcal{A}')$ are compatible if every chart $(U_\alpha, \phi_\alpha) \in \mathcal{A}$ is a CR chart for $(E, \mathcal{A}')$ and vice versa. If a CR atlas $\mathcal{A}$ for E is fixed, we drop the atlas from the notation, understanding that if we refer to E we really refer to $(E, \mathcal{A})$.

We should point out that a given vector bundle E may have many incompatible CR vector bundle structures. For instance, the trivial line bundle $E = M \times \mathbb{C}$ with the trivial CR atlas $\mathcal{A}$ consisting of the single chart (M, ϕ) with $\phi(x, u) = (x, u)$ is compatible with the CR atlas $\mathcal{A}'$ consisting of the single chart (M, ϕ') with $\phi'(x, u) = (x, f(x)u)$, where $f : M \rightarrow \mathbb{C}$ is some nonvanishing function, if and only if f is CR.

2.2. The canonical CR structure. In what follows, $(E, \mathcal{A})$ will denote a CR vector bundle over the CR manifold M . We shall keep the CR atlas $\mathcal{A}$ fixed and simply refer to the CR vector bundle $(E, \mathcal{A})$ as E . In this section, we shall define a CR structure on E . The main point is that the CR structures induced on $\pi^{-1}(U_\alpha)$ by the product CR structures on $U_\alpha \times \mathbb{C}^k$, via the diffeomorphisms ϕ_α , fit together since the transition functions are CR. The CR structure induced on E in this way can also be characterized by a property of sections of E .

Lemma 9. *Assume that at each point $p \in M$ there exists a CR function g with $\bar{L}g(p) \neq 0$ for some CR vector field $\bar{L}$. Then there exists a unique CR structure on E such that a section $\sigma \in \Gamma(U, E)$ on an open subset $U \subset M$ is CR if and only if for each CR chart (V, ϕ) of E the components of $\text{proj}_2 \circ \phi \circ \sigma$ are CR functions on $U \cap V$. Furthermore, relative to this CR structure on E , the projection π is CR, and for each $p \in M$, there exist a neighbourhood U of p and a local basis of CR sections for E over U .*

Before we start with the proof, let us explain some of the notions used in the lemma. M is of finite type if the Lie algebra generated by the CR vector fields and their conjugates, evaluated at $p \in M$, is all of $\mathbb{C}T_p M$. A CR section on U is a section $\sigma \in \Gamma(U, E)$ which is CR as a map $U \rightarrow E$, i.e. $\sigma_* \mathcal{V}(M)_p \subset \mathcal{V}(E)_{\sigma(p)}$ for all $p \in U$. A local basis of CR sections (on U) is a collection of CR sections $\sigma_1, \dots, \sigma_k \in \Gamma(U, E)$ such that $\sigma_1(p), \dots, \sigma_k(p)$ is a basis of E_p for $p \in U$.

We will use that a bundle chart (U_α, ϕ_α) induces a splitting of the tangent space $T_P E$ for $P \in \pi^{-1}(U_\alpha)$: If we write $\phi_\alpha(P) = (p, v)$, then $\mathbb{C}T_P E \cong \mathbb{C}T_{(p,v)}(U_\alpha \times \mathbb{C}^k) \cong \mathbb{C}T_p M \oplus \mathbb{C}T_v \mathbb{C}^k$. The former isomorphism is given by the push forward $(\phi_\alpha)_*$. To make the latter isomorphism explicit, let ι_1 denote the natural inclusion $\mathbb{C}T_p M \hookrightarrow \mathbb{C}T_p M \oplus \mathbb{C}T_v \mathbb{C}^k$ given by $\iota_1(X) = X \oplus 0$, and similarly $\iota_2 : \mathbb{C}T_v \mathbb{C}^k \hookrightarrow \mathbb{C}T_p M \oplus \mathbb{C}T_v \mathbb{C}^k$. A tangent vector $\iota_1 X + \iota_2 Y \in \mathbb{C}T_p M \oplus \mathbb{C}T_v \mathbb{C}^k$ with $X \in \mathbb{C}T_{p_0} M$ and $Y \in \mathbb{C}T_{v_0} \mathbb{C}^k$ acts on the germ of a smooth function $f(p, v)$ at (p_0, v_0) by $(\iota_1 X + \iota_2 Y)f = X(f(p, v_0)) + Y(f(p_0, v))$.

Proof of Lemma 9. We define $\mathcal{V}(E)_P = (\phi_\alpha^{-1})_*(\mathcal{V}(M)_p \oplus T_v^{(0,1)} \mathbb{C}^k)$. We will show that $\mathcal{V}(E)$ defined this way is the unique CR structure claimed in Lemma 9. To start, let us first show that this definition is independent of the chart used. Consider a chart change $\phi = \phi_\alpha \circ \phi_\beta^{-1}$. We denote the fiber coordinate in the chart ϕ_α by $u = (u^1, \dots, u^k)$, the fiber coordinate in the chart ϕ_β by $v = (v^1, \dots, v^k)$, and let $P_0 \in \pi^{-1}(U_\alpha \cap U_\beta)$, $\phi_\alpha(P_0) = (p_0, u_0)$, $\phi_\beta(P_0) = (p_0, v_0)$; thus,

$$(p, u) = \phi(p, v) = (p, \phi_{\alpha\beta}(p)v).$$

We must show that $\phi_*(\mathcal{V}(M)_{p_0} \oplus T_{v_0}^{(0,1)}\mathbb{C}^k) = \mathcal{V}(M)_{p_0} \oplus T_{v_0}^{(0,1)}\mathbb{C}^k$, or equivalently $\phi_*(\iota_1\mathcal{V}(M)_{p_0}) \subset \mathcal{V}(M)_{p_0} \oplus T_{v_0}^{(0,1)}\mathbb{C}^k$ and $\phi_*(\iota_2 T_{v_0}^{(0,1)}\mathbb{C}^k) \subset \mathcal{V}(M)_{p_0} \oplus T_{v_0}^{(0,1)}\mathbb{C}^k$. For this purpose, let $\bar{L}_{p_0} \in \mathcal{V}(M)_{p_0}$. Then

$$\begin{aligned} (\phi_*\iota_1\bar{L}_{p_0})f(p, u) &= \bar{L}_{p_0}f(p, \phi_{\alpha\beta}(p)v_0) \\ &= \bar{L}_{p_0}f(p, \phi_{\alpha\beta}(p_0)v_0) + \frac{\partial f}{\partial u}\bar{L}_{p_0}(\phi_{\alpha\beta}(p))v_0 + \frac{\partial f}{\partial \bar{u}}\bar{L}_{p_0}(\overline{\phi_{\alpha\beta}(p)})\bar{v}_0 \\ &= \iota_1\bar{L}_{p_0}f + \left(\iota_2\frac{\partial}{\partial \bar{u}}\Big|_{u_0} f\right)\bar{L}_{p_0}(\overline{\phi_{\alpha\beta}(p)})\bar{v}_0; \end{aligned}$$

the last equality holds since $\phi_{\alpha\beta}$ is CR. Also,

$$\begin{aligned} (\phi_*\iota_2\frac{\partial}{\partial \bar{v}^j}\Big|_{v_0})f(p, u) &= \iota_2\frac{\partial}{\partial \bar{v}^j}\Big|_{v_0} f(p, \phi_{\alpha\beta}(p)v) \\ &= \frac{\partial}{\partial \bar{v}^j}\Big|_{v_0} f(p_0, \phi_{\alpha\beta}(p_0)v) \\ &= \left(\iota_2\frac{\partial}{\partial \bar{u}}\Big|_{u_0} f\right)\phi_{\alpha\beta}(p_0), \end{aligned}$$

so that $\phi_*\iota_1\bar{L}_{p_0} \in \mathcal{V}(M)_{p_0} \oplus T_{u_0}^{(0,1)}\mathbb{C}^k$ and $\phi_*\iota_2\frac{\partial}{\partial \bar{v}^j}\Big|_{v_0} \in T_{u_0}^{(0,1)}\mathbb{C}^k$. Hence $\phi_*(\mathcal{V}(M)_{p_0} \oplus T_{v_0}^{(0,1)}\mathbb{C}^k) = \mathcal{V}(M)_{p_0} \oplus T_{u_0}^{(0,1)}\mathbb{C}^k$, as claimed.

It is easy to check that $[\mathcal{V}(E), \mathcal{V}(E)] \subset \mathcal{V}(E)$ (since the derivatives in the fiber directions and the derivatives in the base directions commute). It also follows immediately that $\mathcal{V}(E) \cap \overline{\mathcal{V}(E)} = \{0\}$. We conclude that $\mathcal{V}(E)$ defines a CR structure on E ; furthermore, $\pi_*(\mathcal{V}(E)) = \mathcal{V}(M)$, so that π is CR as claimed.

We next verify that $\mathcal{V}(E)$ has the properties claimed in Lemma 9. Let $\sigma : U \rightarrow E|_U$ be a section, and (V, ϕ) a CR chart with $U \cap V$ nonempty. By the construction of the CR structure on E , σ is CR if and only if $\phi \circ \sigma$ is CR. Since σ is a section, we have $\phi \circ \sigma(p) = (p, \tilde{\sigma}(p))$, where $\tilde{\sigma} = \text{proj}_2 \circ \phi \circ \sigma = (\sigma^1, \dots, \sigma^k)^t$. We compute $(\phi \circ \sigma)_*\bar{L}_{p_0}$, where, for ease of notation, we drop the inclusions ι_1, ι_2 :

$$\begin{aligned} ((\phi \circ \sigma)_*\bar{L}_{p_0})f(p, u) &= \bar{L}_{p_0}f(p, \tilde{\sigma}(p)) \\ &= \bar{L}_{p_0}f(p, \tilde{\sigma}(p_0)) + \frac{\partial f(p_0, u)}{\partial \bar{u}}\Big|_{\tilde{\sigma}(p_0)}\bar{L}_{p_0}\tilde{\sigma} + \frac{\partial f(p_0, u)}{\partial u}\Big|_{\tilde{\sigma}(p_0)}\bar{L}_{p_0}\tilde{\sigma}. \end{aligned}$$

From this we see that $\phi \circ \sigma$ (and hence σ) is CR if and only if the components of $\tilde{\sigma}$ are CR functions.

Now let $p \in M$ and assume that there exists a CR function g with the property that $\bar{L}g(p) \neq 0$. Let us show that the CR structure constructed above is unique with the properties in Lemma 9. Pick $P_0 \in E$ and a CR chart (V, ϕ) with $P_0 \in \pi^{-1}(V)$. Write $\phi(P_0) = (p_0, u_0)$. There is no loss of generality in assuming that $g(p_0) = u_0$. Using the computation above, first with $\tilde{\sigma} = u_0$ and then with $\tilde{\sigma} = ge_j$ for standard unit vectors e_j , $j = 1, \dots, k$, we conclude, since $\sigma = \phi^{-1}(p, \tilde{\sigma})$ is CR for each of the choices of $\tilde{\sigma}$ above, that $\mathcal{V}(E)$ must be the bundle $(\phi^{-1})_*(\mathcal{V}(M)_{p_0} \oplus T_{u_0}^{(0,1)}\mathbb{C}^k)$. This completes the proof of the Lemma. $\square$

Remark 1. In what follows, unless specifically stated otherwise, every CR vector bundle will be equipped with the canonical CR structure provided by Lemma 9.

2.3. Constructions with CR vector bundles. All standard vector bundle constructions work in the category of CR vector bundles. In particular, the sum, tensor product, and exterior product of CR vector bundles carry the structure of a CR vector bundle, and the dual bundle E^* is a CR vector bundle. Also, if $\hat{E}$ is a CR vector bundle over another CR manifold $\hat{M}$, and $f : M \rightarrow \hat{M}$ is a CR map, the pullback bundle $f^*\hat{E}$ inherits the structure of a CR vector bundle.

Let us discuss the construction of the pullback bundle in detail. We let $E_x = \hat{E}_{f(x)}$ and define a vector bundle structure on $E = \cup_{x \in M} E_x$ as follows: If $\hat{U} \subset \hat{M}$ is open and $\hat{\phi} : (\hat{\pi})^{-1}(\hat{U}) \rightarrow \hat{U} \times \mathbb{C}^k$ is a CR chart for $\hat{E}$ over $\hat{U}$, $(\hat{\phi}^{-1} \circ (f \times \text{id}))^{-1}$ is a bundle chart for E over $U = f^{-1}(\hat{U})$. Since the transition functions for E are just the pullbacks (by f) of the transition functions for $\hat{E}$, E is a CR vector bundle. The map $f : M \rightarrow \hat{M}$ lifts to a differentiable vector bundle map $F : E \rightarrow \hat{E}$. The canonical CR structure on $\hat{E}$ is connected with the structure on E by $\mathcal{V}(E)_p \subset \mathcal{V}(\hat{E})_{f(p)}$, so that $F : E \rightarrow \hat{E}$ is a CR map. Also note that the pullback (by f) $f^*\sigma \in \Gamma(f^{-1}(\hat{U}), E)$ of a CR section $\sigma \in \Gamma(\hat{U}, \hat{E})$ is a CR section of E .

2.4. The derivative operator. In analogy with complex analytic vector bundles, we can differentiate sections of a CR vector bundle E (equipped with the canonical CR structure provided by Lemma 9) with respect to CR vector fields. To see this, let $\sigma_1, \dots, \sigma_k$ be a local basis of CR sections over some open set U . If $\sigma \in \Gamma(U, E)$ is given by $\sigma = \sum_{j=1}^k a^j \sigma_j$ with $a^j \in C^\infty(U)$ and $\bar{L}$ is a CR vector field on U , we define

$$D_{\bar{L}}\sigma = \sum_{j=1}^k (\bar{L}a^j) \sigma_j.$$

This definition is invariant under changes of the local CR basis σ_j : Observe that a section $\sigma \in \Gamma(U, E)$, $\sigma = \sum_{j=1}^k a^j \sigma_j$ is CR if and only if all the a^j are CR functions on U . Hence, if $\tilde{\sigma}_1, \dots, \tilde{\sigma}_k$ is another basis of CR sections, $\sigma_j = \sum_l c_j^l \tilde{\sigma}_l$ where the c_j^l are CR functions on U . It follows that $\sigma = \sum_l (\sum_j c_j^l a^j) \tilde{\sigma}_l = \sum_l \tilde{a}^l \tilde{\sigma}_l$, and

$$\begin{aligned} \sum_l (\bar{L}\tilde{a}^l) \tilde{\sigma}_l &= \sum_l \bar{L}(\sum_j c_j^l a^j) \tilde{\sigma}_l \\ &= \sum_l (\sum_j c_j^l \bar{L}a^j) \tilde{\sigma}_l \\ &= \sum_j (\bar{L}a^j) \sigma_j, \end{aligned}$$

so that $D_{\bar{L}}\sigma$ is independent of the local basis of CR sections used. This allows us to define $D_{\bar{L}}\sigma$ for a section $\sigma \in \Gamma(U, E)$ on any open set $U \in M$. Note that σ is CR if and only if $D_{\bar{L}}\sigma = 0$ (the 0 section) for all CR vector fields $\bar{L}$.

We also note that the operator $\omega \mapsto D_{\bar{L}}\omega$ is local, and since $D_{a\bar{L}}\omega = aD_{\bar{L}}\omega$ for any function a , $(D_{\bar{L}}\omega)_p$ depends only on $\bar{L}_p$. Moreover, D satisfies the product rule:

$$(7) \quad D_{\bar{L}}(a\omega) = (\bar{L}a)\omega + aD_{\bar{L}}\omega.$$

If $\omega = (\omega_1, \dots, \omega_k)$ is any local (not necessarily CR) basis of sections over U and $\bar{L}_1, \dots, \bar{L}_n$ is a local basis of CR vector fields over U , we can write $D_{\bar{L}_i}\omega_j = \sum_l a_{ij}^l \omega_l$, (or, using matrix notation, $D_{\bar{L}_i}\omega = A_i\omega$) and $A_i = (a_{ij}^l)_i$ determines $D_{\bar{L}}\sigma$ for all sections σ over U . If we change basis by $\tilde{\omega}_j = \sum_m C_j^m \omega_m$, which

we will write in matrix notation as $\tilde{\omega} = C\omega$, we obtain

$$\begin{aligned} D_{\bar{L}_i} \tilde{\omega} &= D_{\bar{L}_i} C\omega \\ &= (\bar{L}_i C)\omega + C(D_{\bar{L}_i} \omega) \\ &= ((\bar{L}_i C)C^{-1} + CA_i C^{-1})\tilde{\omega}. \end{aligned}$$

This means that under a change of basis, the A_i transform as follows:

$$\tilde{\omega} = C\omega \implies \tilde{A}_i = (\bar{L}_i C)C^{-1} + CA_i C^{-1}.$$

We can also think of D as an operator from sections of E to sections of $E \otimes \mathcal{V}(M)^* \simeq E \otimes (\mathbb{C}T^*M/\mathcal{V}(M)^\perp)$. Since D satisfies the product rule (7), we may consider D as a partial connection in E .

If E is the trivial bundle $M \times \mathbb{C}$ or the *holomorphic cotangent bundle* $T'M = \mathcal{V}(M)^\perp$, then D is just $\bar{\partial}_b$. If M is a complex manifold and E is an analytic vector bundle, D is $\bar{\partial}$.

3. THE CR VECTOR BUNDLE STRUCTURE ON THE HOLOMORPHIC COTANGENT BUNDLE

3.1. Definition. Let now M be an *integrable* CR manifold of CR dimension $n = \dim \mathcal{V}(M)_p$ and CR codimension $d = \dim M - 2n$. Integrability means that at each point $p \in M$ there exist a neighbourhood U of p and $N = n + d$ smooth CR functions Z_j on U , $j = 1, \dots, N$ such that the differentials $dZ_1, \dots, dZ_N$ are linearly independent on U . The functions $Z_1, \dots, Z_N$ are called a *family of basic solutions* in U . One can show that $q \mapsto (Z_1(q), \dots, Z_N(q))$ embeds M , locally near $p \in M$, as a generic submanifold in $\mathbb{C}^N$. Conversely, any embeddable (in this sense) CR manifold is integrable.

We recall the definition of the holomorphic cotangent bundle $T'M$: At each $p \in M$, $T'_p M = \mathcal{V}(M)_p^\perp$. Since $\dim \mathbb{C}T_p M = 2n + d$ and $\dim \mathcal{V}(M)_p = n$, $\dim T'_p M = N$. Hence, on a neighbourhood U of a fixed point p as above, the 1-forms $dZ_1, \dots, dZ_N$ form a basis of the sections of $T'M$ over U . We want to define a CR atlas on $T'M$ such that the bases obtained in this way are CR bases of sections. Assume that we have another family of basic solutions $\tilde{Z}_1, \dots, \tilde{Z}_N$ on the open set $V \in M$. Then $d\tilde{Z}_j = \sum_k a_j^k dZ_k$ on $U \cap V$, and exterior differentiation yields

$$\sum_k da_j^k \wedge dZ_k = 0$$

in the overlap $U \cap V$. By Cartan's Lemma, we conclude that $da_j^k = \sum_l C_k^l dZ_l$, and hence, $\bar{L}a_j^k = \langle da_j^k, \bar{L} \rangle = 0$ for all CR vector fields $\bar{L}$, so that the a_j^k are CR functions on $U \cap V$. Hence, if we define a CR chart for $T'M$ over U by $\phi(\omega_p) = (p, \lambda^1, \dots, \lambda^N)$ where $\omega_p \in T'_p M$ is written as $\omega_p = \sum_j \lambda^j (dZ_j)_p$, then, by the computation above, the collection of all such charts defines a CR atlas on $T'M$. We equip $T'M$ with the canonical CR structure, relative to this CR atlas, given by Lemma 9.

By the same argument as above using Cartan's lemma, we see that the following lemma holds.

Lemma 10. *Let g be a smooth function on an open set $U \subset M$. Then g is CR if and only if dg is a CR section of $T'M$ over U .*

Sections of $T'M$ are called *holomorphic forms* (on M). Let now $\omega \in \Gamma(M, T'M)$, and let $\bar{L}$ be a CR vector field on M . The following useful formula holds:

$$(8) \quad D_{\bar{L}} \omega = \bar{L} \lrcorner d\omega.$$

To verify this identity, let $X \in \Gamma(M, \mathbb{C}TM)$ be a tangent vector field, and $\omega = \sum_j a^j dZ_j$ locally. Then

$$\begin{aligned} \langle \bar{L} \lrcorner d\omega, X \rangle &= \langle d\omega, \bar{L} \wedge X \rangle \\ &= \sum_j \langle da^j \wedge dZ_j, \bar{L} \wedge X \rangle \\ &= \sum_j (da^j(\bar{L}) dZ_j(X) - da^j(X) dZ_j(\bar{L})) \\ &= \sum_j (\bar{L}a^j) dZ_j(X) \\ &= \langle D_{\bar{L}}\omega, X \rangle. \end{aligned}$$

3.2. Nondegeneracy conditions. Recall that the *characteristic bundle* T^0M is defined by

$$T^0M = \left(\mathcal{V}(M) \oplus \overline{\mathcal{V}(M)} \right)^\perp.$$

A *characteristic form* is a real nonvanishing section of T^0M , and we write $\xi^0(U)$ for the space of characteristic forms on $U \subset M$. We usually denote a characteristic form by θ .

By the discussion above, an integrable CR manifold M is finitely nondegenerate (at p) as introduced by Baouendi, Huang and Rothschild [3] if and only if for some k ,

$$\text{span}\{(D_{\bar{L}_1} \dots D_{\bar{L}_l} \theta)(p) : \bar{L}_j \in \Gamma(U, \mathcal{V}(M)), \theta \in \xi^0(U), l \leq k\} = T'_p M,$$

where U is a suitable small neighbourhood of p . This follows from the characterization of finite nondegeneracy found in e.g. [2] and equation (8).

We will use the notion of the pullback bundle introduced before in order to give an intrinsic characterization of the nondegeneracy of a CR map $f : M \rightarrow \hat{M}$ as introduced in [9]. Let $E = f^*T'\hat{M}$. f is nondegenerate at p_0 if and only if for some k ,

$$\text{span}\{D_{\bar{L}_1} \dots D_{\bar{L}_k} f^\# \theta(p_0) : \bar{L}_j \in \Gamma(U, \mathcal{V}(M)), \theta \in \hat{\xi}(\hat{U}), j \leq k\} = E_{p_0},$$

where again U is some neighbourhood of p_0 , contained in $f^{-1}(\hat{U})$. More generally, we can define subspaces $E_p^k \subset E_p$ (for $p \in U$, say) by

$$E_p^k = \text{span}\{D_{\bar{L}_1} \dots D_{\bar{L}_l} \theta(p) : \bar{L}_j \in \Gamma(U, \mathcal{V}(M)), \theta \in \xi'(U), l \leq k\}$$

Then $E_p^0 \subset E_p^1 \subset \dots \subset E_p^k \subset \dots$, and for some smallest integer $k_0 = k_0(p)$, $E_p^k = E_p^{k_0}$ for $k \geq k_0$. We call $s = s(p) = \dim E_p - \dim E_p^{k_0(p)}$ the degeneracy of f at p and say that f is (k_0, s) -degenerate at p ; we say that f is of constant degeneracy s at p_0 if $s(p) \equiv d$ is constant in a neighbourhood of p_0 , and in this case we also say that f is constantly (k_0, s) -degenerate (note that the constancy only applies to s). This characterization is equivalent to the definition given in [8], by a computation in coordinates which we leave to the reader.

Note that $p \mapsto s(p)$ is upper semicontinuous. Hence, we have the following:

Lemma 11. *Let $f : M \rightarrow \hat{M}$ be a smooth CR map. Then there exists an open dense subset $N \subset M$ such that f is of constant degeneracy at every point of N .*

Let us actually observe some more. It is proved in [8] that the degeneracy s of a transversal map is bounded by $\hat{N} - N$. Furthermore it follows from that paper (and is easy to prove), that $\dim E_p^1 = N$ for any point p , if $\hat{M}$ is Levi nondegenerate and f is an embedding. Now note that if on any open subset U we had $E_p^{k+1} \subset E_p^k$ for $p \in U$, it follows that $E_p = E_p^k$. It follows that if f is of constant degeneracy s on U , then generically in U , $k_0 \leq \hat{N} - N + 1 - s \leq \hat{N} - N + 1$. We state this as a Lemma:

Lemma 12. *Let $f: M \rightarrow \hat{M}$ be a smooth transversal CR embedding between Levi nondegenerate hypersurfaces M and $\hat{M}$. Then there exists an open dense subset $N \subset M$ such that f is of constant degeneracy (k_0, s) at every point of N for some $s \leq \hat{N} - N$ and $k_0 \leq \hat{N} - N + 1 - s \leq \hat{N} - N + 1$.*

3.3. Some computations. From now on, unless stated otherwise, we assume that M and $\hat{M}$ are of hypersurface type and integrable (so that the holomorphic cotangent bundles are CR vector bundles), and that $f: M \rightarrow \hat{M}$ is CR. Let us now fix a point p_0 (which we denote by 0 from now on) and a local basis $L_{\bar{1}}, \dots, L_{\bar{n}}$ of the CR vector fields on M near 0. We also choose a characteristic form θ for M near 0 and a real vector field T with $\langle \theta, T \rangle = 1$. We denote by θ^a the dual form (relative to the basis of vector fields $T, L_1, \dots, L_n, L_{\bar{1}}, \dots, L_{\bar{n}}$) to the anti CR vector field L_a . We write $\hat{0} = f(0)$ and, as before, $\hat{\theta}$ denotes a (local) characteristic form on $\hat{M}$ near $\hat{0}$, and $\hat{T}$ a real vector field with $\langle \hat{\theta}, \hat{T} \rangle = 1$; we extend $\hat{\theta}$ to a basis of $T'\hat{M}$ by $\hat{\theta}^1, \dots, \hat{\theta}^{\hat{n}}$ which are the dual forms of a local basis $\hat{L}_1, \dots, \hat{L}_{\hat{n}}$ of the anti CR vector fields on $\hat{M}$. We work on a small neighbourhood of 0 which we may shrink, if necessary, without further mentioning it.

Since f is CR, we can write

$$f_* L_a = \gamma_a^A \hat{L}_A, \quad f_* L_{\bar{a}} = \overline{\gamma_a^A} \hat{L}_{\bar{A}}, \quad f_* T = \eta^A \hat{L}_A + \overline{\eta^A} \hat{L}_{\bar{A}} + \xi \hat{T},$$

or equivalently,

$$f^* \hat{\theta}^A = \gamma_a^A \theta^a + \eta^A \theta, \quad f^* \hat{\theta}^{\bar{A}} = \overline{\gamma_a^A} \theta^{\bar{a}} + \overline{\eta^A} \theta, \quad f^* \hat{\theta} = \xi \theta.$$

Here and from now on we use the summation convention; small indices a, b etc. range from 1 to n , capital indices A, B etc. from 1 to $\hat{n}$. We observe here that f is transversal at $p \in M$, as defined in the introduction, if and only if $\xi(p) \neq 0$.

Let us write $D_{\bar{a}} = D_{L_{\bar{a}}}$, $f^{\#} \hat{\theta} = \tau$, and $f^{\#} \hat{\theta}^A = \tau^A$; thus, $\tau, \tau^1, \dots, \tau^n$ forms a local basis (not necessarily CR) for the CR vector bundle $E = f^{\#} T' \hat{M}$. We define functions $g_{\bar{a}_1 \dots \bar{a}_k B}$, $g_{\bar{a}_1 \dots \bar{a}_k}$ on M near 0 by

$$D_{\bar{a}_k} D_{\bar{a}_{k-1}} \dots D_{\bar{a}_1} \tau = g_{\bar{a}_1 \dots \bar{a}_k B} \tau^B + g_{\bar{a}_1 \dots \bar{a}_k} \tau.$$

Note that for elements σ_p of E_p , there is a natural pairing $\langle \sigma_p, X_{f(p)} \rangle$ with elements $X_{f(p)}$ of $\mathbb{C} T_{f(p)} \hat{M}$. With this notation, we can write

$$g_{\bar{a}_1 \dots \bar{a}_k B} = \langle D_{\bar{a}_k} D_{\bar{a}_{k-1}} \dots D_{\bar{a}_1} \tau, \hat{L}_B \rangle, \quad g_{\bar{a}_1 \dots \bar{a}_k} = \langle D_{\bar{a}_k} D_{\bar{a}_{k-1}} \dots D_{\bar{a}_1} \tau, \hat{T} \rangle.$$

We introduce similar notation for sections of $T' \hat{M}$ and $T' M$:

$$D_{\bar{a}_k} \dots D_{\bar{a}_1} \theta = h_{\bar{a}_1 \dots \bar{a}_k b} \theta^b + h_{\bar{a}_1 \dots \bar{a}_k} \theta,$$

$$D_{\bar{A}_k} \dots D_{\bar{A}_1} \hat{\theta} = \hat{h}_{\bar{A}_1 \dots \bar{A}_k B} \hat{\theta}^B + \hat{h}_{\bar{A}_1 \dots \bar{A}_k} \hat{\theta}.$$

With $\langle \cdot, \cdot \rangle$ denoting the canonical pairing between vectors and covectors,

$$h_{\bar{a}_1 \dots \bar{a}_k b} = \langle D_{\bar{a}_k} D_{\bar{a}_{k-1}} \dots D_{\bar{a}_1} \theta, L_b \rangle, \quad h_{\bar{a}_1 \dots \bar{a}_k} = \langle D_{\bar{a}_k} D_{\bar{a}_{k-1}} \dots D_{\bar{a}_1} \theta, T \rangle,$$

$$\hat{h}_{\bar{A}_1 \dots \bar{A}_k B} = \langle D_{\bar{A}_k} D_{\bar{A}_{k-1}} \dots D_{\bar{A}_1} \hat{\theta}, \hat{L}_B \rangle, \quad \hat{h}_{\bar{A}_1 \dots \bar{A}_k} = \langle D_{\bar{A}_k} D_{\bar{A}_{k-1}} \dots D_{\bar{A}_1} \hat{\theta}, \hat{T} \rangle.$$

The $h_{\bar{a}b}$ relate to the Leviform of M by

$$(9) \quad h_{\bar{a}b} = \langle d\theta, L_{\bar{a}} \wedge L_b \rangle = -\theta([L_{\bar{a}}, L_b]),$$

and similarly for $\hat{M}$.

We need to introduce (but just for a short while) one more notation for our three bundles:

$$D_{\bar{a}} \theta^b = R_{\bar{a}c}^b \theta^c + R_{\bar{a}}^b \theta,$$

$$D_{\bar{a}} \tau^B = S_{\bar{a}C}^B \tau^C + S_{\bar{a}}^B \tau,$$

$$D_{\bar{A}}\hat{\theta}^B = \hat{R}_{\bar{A}C}^B \hat{\theta}^C + \hat{R}_{\bar{A}}^B \hat{\theta}.$$

Lemma 13. *Let σ be a section of $T'\hat{M}$. Then*

$$(10) \quad D_{\bar{a}}f^\sharp\sigma = \overline{\gamma_a^A}f^\sharp D_{\bar{A}}\sigma.$$

In particular,

$$(11) \quad g_{\bar{a}B} = \overline{\gamma_a^A}\hat{h}_{\bar{A}B}, \quad g_{\bar{a}} = \overline{\gamma_a^A}\hat{h}_{\bar{A}},$$

and

$$(12) \quad S_{\bar{a}C}^B = \overline{\gamma_a^A}\hat{R}_{\bar{A}C}^B, \quad S_{\bar{a}}^C = \overline{\gamma_a^A}\hat{R}_{\bar{A}}^C.$$

Proof. Let us consider a local CR basis $\omega_1, \dots, \omega_{\hat{N}}$ for $T'\hat{M}$. We write $\alpha_j = f^\sharp\omega_j$ for the corresponding local CR basis of E . Then locally $\sigma = a^j\omega_j$ for smooth functions a^j , and we have

$$\begin{aligned} D_{\bar{a}}f^\sharp\sigma &= D_{\bar{a}}f^\sharp(a^j\omega_j) \\ &= D_{\bar{a}}(a^j \circ f)\alpha_j \\ &= (L_{\bar{a}}(a^j \circ f))\alpha_j \\ &= ((f_*L_{\bar{a}})a^j)\alpha_j \\ &= \overline{\gamma_a^A}(L_{\bar{A}}a^j \circ f)\alpha_j \\ &= \overline{\gamma_a^A}f^\sharp D_{\bar{A}}\sigma. \end{aligned}$$

For the other assertions, we note that

$$D_{\bar{a}}\tau = g_{\bar{a}B}\tau^B + g_{\bar{a}}\tau$$

and

$$D_{\bar{A}}\hat{\theta} = \hat{h}_{\bar{A}B}\hat{\theta}^B + h_{\bar{A}}\hat{\theta}.$$

Since

$$\overline{\gamma_a^A}f^\sharp(D_{\bar{A}}\hat{\theta}) = \overline{\gamma_a^A}\hat{h}_{\bar{A}B}\tau^B + \overline{\gamma_a^A}\hat{h}_{\bar{A}}\tau,$$

comparing coefficients we get (11). (12) follows by the same argument. $\square$

The following lemma is an immediate consequence of (7) and the definitions above:

Lemma 14. *With the notations introduced above, the following hold:*

$$(13) \quad g_{\bar{a}_1\dots\bar{a}_k\bar{a}B} = L_{\bar{a}}g_{\bar{a}_1\dots\bar{a}_k B} + g_{\bar{a}_1\dots\bar{a}_k}S_{\bar{a}B}^C + g_{\bar{a}_1\dots\bar{a}_k}g_{\bar{a}B}$$

$$(14) \quad \hat{h}_{\bar{A}_1\dots\bar{A}_k\bar{A}B} = L_{\bar{A}}\hat{h}_{\bar{A}_1\dots\bar{A}_k B} + \hat{h}_{\bar{A}_1\dots\bar{A}_k} \hat{R}_{\bar{A}B}^C + \hat{h}_{\bar{A}_1\dots\bar{A}_k}\hat{h}_{\bar{A}B}$$

$$(15) \quad h_{\bar{a}_1\dots\bar{a}_k\bar{a}b} = L_{\bar{a}}h_{\bar{a}_1\dots\bar{a}_k b} + h_{\bar{a}_1\dots\bar{a}_k}R_{\bar{a}b}^c + h_{\bar{a}_1\dots\bar{a}_k}h_{\bar{a}b}.$$

We also record the following two lemmas for later use. We write $I = (i_1 \dots i_l)$, where $1 \leq i_j \leq n$, and we use the notation $L^I = L_{i_1} \dots L_{i_l}$; for such an I , we write $|I| = l$.

Lemma 15. *For each $k \in \mathbb{N}$, the two subspaces of $\mathbb{C}^{\hat{n}}$ defined by*

$$\text{span}\{(g_{\bar{a}_1\dots\bar{a}_l A}(p))_A : l \leq k\}$$

$$\text{span}\{(L^I g_{\bar{a}A}(p))_A : |I| \leq k\}$$

coincide. Consequently, the following are equivalent definitions for the degeneracy $d(p)$ of f at p defined in section 3.2:

(i) $d(p) = \hat{n} - \max_k \dim \text{span}\{(g_{\bar{a}_1 \dots \bar{a}_k A}(p))_A : l \leq k\};$

(ii) $d(p) = \hat{n} - \max_k \dim \text{span}\{(L^I g_{\bar{a} A}(p))_A : |I| \leq k\};$

and in each case, $k_0(p)$ is the minimal integer k for which the maximum on the right hand side is attained.

The proof is by induction using (13).

Lemma 16. *For each $a_1, \dots, a_k$, $1 \leq a_l \leq n$, and B , $1 \leq B \leq \hat{n}$ there exist functions $u_{\bar{a}_1 \dots \bar{a}_k B}$ and $u_{\bar{a}_1 \dots \bar{a}_k}$ polynomial in their arguments with coefficients smooth on $M \times \hat{M}$ such that*

$$(16) \quad g_{\bar{a}_1 \dots \bar{a}_k B} = u_{\bar{a}_1 \dots \bar{a}_k B} \left(\overline{L^I \gamma_a^A} \right), \quad |I| \leq k-1$$

$$(17) \quad g_{\bar{a}_1 \dots \bar{a}_k} = u_{\bar{a}_1 \dots \bar{a}_k} \left(\overline{L^I \gamma_a^A} \right), \quad |I| \leq k-1.$$

Furthermore, $u_{\bar{a}_1 \dots \bar{a}_k A}$ and $u_{\bar{a}_1 \dots \bar{a}_k}$ depend only on M and $\hat{M}$ (and not on the mapping f).

The equalities (16) and (17) mean the following: if the left hand side is evaluated at p , it equals the right hand side with the coefficients evaluated at $(p, f(p))$ and the arguments evaluated at p . We will encounter many such equalities, and we will always use this time-saving notation.

For the proof of Lemma 16, we recall that for $k=1$, $g_{\bar{a} B} = \overline{\gamma_a^A} \hat{h}_{\bar{A} B}$ and $g_{\bar{a}} = \overline{\gamma_a^A} \hat{h}_{\bar{A}}$ by Lemma 13. An induction and Lemma 14 finishes the proof of the Lemma 16; the details are left to the reader.

3.4. The pullback operation. In this section, we discuss the relation between our vector bundle $E = f^\# T' \hat{M}$ and the vector bundle $T' M$ over M . Recall that we have a pairing $\langle \sigma, X \rangle$ for sections of E and tangent vectors X at $f(p)$. We define a map ϕ by

$$\langle \phi \sigma, X \rangle = \langle \sigma, f_* X \rangle.$$

Since f is CR, ϕ maps sections of E to sections of $T' M$. Note that $\phi : \Gamma(U, E) \rightarrow \Gamma(U, T' M)$ is linear over $C^\infty(U)$, since

$$\langle \phi a \sigma, X \rangle = (a \sigma)(f_* X) = a \sigma(f_* X) = a \langle \phi \sigma, X \rangle.$$

Lemma 17. *For each j , $D_{\bar{j}} \phi = \phi D_{\bar{j}}$. Furthermore, $\phi \circ f^\# = f^*$.*

Proof. Since the derivative operation is local, it is actually enough to work on a trivializing open set $U = f^{-1}(V)$, where we have a family of basic solutions $\hat{Z}_j$ in V . A basis of the CR sections of E is then given by $\sigma_j = f^\# d\hat{Z}_j$. Let τ be any section of E over U . Then $\tau = a^j \sigma_j$, and $\phi \tau = a^j \phi(\sigma_j)$. By the definition of $D_{\bar{j}}$ we now see it is enough to show that if σ is a CR section of E , then $\phi \sigma$ is a CR section of $T' M$. We compute:

$$\begin{aligned} \langle (\phi f^\# d\hat{Z}_j)_p, X_p \rangle &= \langle (f^\# d\hat{Z}_j)_p, f_* X_p \rangle \\ &= \langle (d\hat{Z}_j)_{f(p)}, f_* X_p \rangle \\ &= \langle f^*(d\hat{Z}_j)_p, X_p \rangle \\ &= \langle d(\hat{Z}_j \circ f)_p, X_p \rangle, \end{aligned}$$

so that actually $\phi f^\# d\hat{Z}_j = d(\hat{Z}_j \circ f)$, which is CR by Lemma 10. The second claim follows from the same calculation. $\square$

By an appropriate choice of basis θ^a , θ , and similarly on $\hat{M}$, we may assume that $R_{\bar{a}c}^b, R_{\bar{a}}^b, \hat{R}_{AC}^B, R_A^B$ are all 0 (cf. [5], Lemma 1.33). By Lemma 13, it follows that $S_{\bar{a}c}^B = S_{\bar{a}}^B = 0$. Lemma 17 implies the following relations:

$$(18) \quad \xi h_{\bar{a}b} = \gamma_a^B g_{\bar{b}B};$$

$$(19) \quad L_{\bar{a}}\xi + \xi h_{\bar{a}} = g_{\bar{a}B}\eta^B + g_{\bar{a}}\xi.$$

This follows comparing the coefficients of $D_{\bar{a}}\phi\tau$ and $\phi D_{\bar{a}}\tau$. We also need the following equalities:

$$(20) \quad L_{\bar{a}}\gamma_b^B = -\eta^B h_{\bar{a}b},$$

$$(21) \quad L_{\bar{a}}\eta^B = -\eta^B h_{\bar{a}},$$

$$(22) \quad T\gamma_a^A = L_a\eta^A + \eta^A h_a,$$

which follow as in [5]. We observe that the identity (18) implies that if f is transversal at p (i.e. $\xi(p) \neq 0$) and M is Levi nondegenerate at p (i.e. the matrix $(h_{\bar{a}b})(p)$ is invertible), then the rank of the matrix $(\gamma_a^B)(p)$ must be maximal, i.e. $= n$. Thus, if f is transversal and M is Levi nondegenerate, then f is in fact a local embedding.

4. JET REFLECTION IDENTITIES FOR NONDEGENERATE MAPPINGS

In this section we are going to establish the jet reflection identities necessary to use the machinery developed in [5]. Our first goal is the reflection identities for nondegenerate mappings. Fix an integer k_0 . We denote by $\mathcal{K}$ the set of all strings $\mathbb{A} = \bar{a}_1 \dots \bar{a}_k$ of elements of $\{\bar{1}, \dots, \bar{n}\}$ with $k \leq k_0$, and we write $\mathcal{K}_{\hat{n}}$ for the set of all $\hat{n}$ -tuples of elements of $\mathcal{K}$. For $\mathbb{A} = \bar{a}_1 \dots \bar{a}_k$ we let $\ell(\mathbb{A}) = k$ denote the length of $\mathbb{A}$.

Theorem 18. *Let $k_0 \in \mathbb{N}$. There exist functions*

$$\Delta_{\mathbf{A}} \left(\overline{L^J \gamma_c^D} \right), \quad r_{a,\mathbf{A}}^B \left(\overline{L^J \gamma_c^D}, \overline{L^I \xi} \right), \quad r_{\mathbf{A}}^B \left(\overline{L^J \gamma_c^D}, \overline{L^I \xi} \right),$$

where $|J| \leq k_0 - 1, |I| \leq k_0$, for $\mathbf{A} = (\mathbf{A}_0, \dots, \mathbf{A}_n) \in \mathcal{K}_{\hat{n}}^{n+1}$, $a \in \{1, \dots, n\}$ and $B \in \{1, \dots, \hat{n}\}$ with the following property: If $f : M \rightarrow \hat{M}$ is k -nondegenerate at 0 with $k \leq k_0$, then there exists $\mathbf{A}^0$ with $\Delta_{\mathbf{A}^0} \left(\overline{L^J \gamma_c^D}(0) \right) \neq 0$, and for any $\mathbf{A} \in \mathcal{K}_{\hat{n}}^{n+1}$ with $\Delta_{\mathbf{A}} \left(\overline{L^J \gamma_c^D}(0) \right) \neq 0$ the following equalities hold:

$$(23) \quad \gamma_a^B = r_{a,\mathbf{A}}^B \left(\overline{L^J \gamma_c^D}, \overline{L^I \xi} \right),$$

$$(24) \quad \eta^B = r_{\mathbf{A}}^B \left(\overline{L^J \gamma_c^D}, \overline{L^I \xi} \right).$$

The $\Delta_{\mathbf{A}}$ are polynomials in $\overline{L^J \gamma_c^D}$ with complex coefficients; the functions $r_{a,\mathbf{A}}^B$ and $r_{\mathbf{A}}^B$ are rational in $\overline{L^J \gamma_c^D}$, polynomial in $\overline{L^I \xi}$, with coefficients smooth on $M \times \hat{M}$. Furthermore, $\Delta_{\mathbf{A}}$, $r_{a,\mathbf{A}}^B$, and $r_{\mathbf{A}}^B$ only depend on M , $\hat{M}$, and k_0 (and not on the mapping f).

In order to define the functions $\Delta_{\mathbf{A}}$ of the Theorem, we need the following computational Lemma.

Lemma 19. *Let $f : M \rightarrow \hat{M}$ be a CR mapping. Then for each $k \in \mathbb{N}$, all $a \in \{1, \dots, n\}$, and all $\mathbb{A} = \bar{a}_1 \dots \bar{a}_k$ the following identities hold:*

$$(25) \quad g_{\mathbb{A}B}\eta^B = \sum_{\ell(\mathbb{A}') < \ell(\mathbb{A})} g_{\mathbb{A}'B}\eta^B v_{\mathbb{A}}^{\mathbb{A}'} \left(\overline{L^J \gamma_c^D} \right) + w_{\mathbb{A}} \left(\overline{L^J \gamma_c^D}, \overline{L^I \xi} \right);$$

$$(26) \quad g_{\mathbb{A}B}\gamma_a^B = \sum_{\ell(\mathbb{A}') < \ell(\mathbb{A})} g_{\mathbb{A}'B}\gamma_a^B p_{a\mathbb{A}}^{\mathbb{A}'} \left(\overline{L^J \gamma_c^D} \right) + \sum_{\ell(\mathbb{A}') < \ell(\mathbb{A})} g_{\mathbb{A}'} \eta^B q_{a\mathbb{A}}^{\mathbb{A}'} \left(\overline{L^J \gamma_c^D} \right) + r_{a\mathbb{A}} \left(\overline{L^J \gamma_c^D}, \overline{L^I \xi} \right);$$

where $|J| \leq k-1$, $|I| \leq k$. The functions v , w , p , q , and r are rational in $\overline{L^I \gamma_c^D}$, the functions w and r are polynomial in $\overline{L^I \xi}$, with coefficients smooth on $M \times \hat{M}$. Furthermore, v , w , p , q , and r only depend on M and $\hat{M}$.

Proof. We start with (25). For $k=1$, by (19),

$$g_{\bar{a}B}\eta^B = L_{\bar{a}}\xi + \xi h_{\bar{a}} - \xi g_{\bar{a}},$$

and recalling (11), $g_{\bar{a}} = \overline{\gamma_a^A} h_{\mathbb{A}}$, we have that

$$g_{\bar{a}B}\eta^B = L_{\bar{a}}\xi + \xi h_{\bar{a}} - \xi \overline{\gamma_a^A} h_{\mathbb{A}} = w_{\bar{a}} \left(\overline{\gamma_c^D}, \overline{L^I \xi} \right),$$

which is (25) for $k=1$. To get (25) for $k>1$, we use (25) for $k-1$ to express $g_{\bar{a}_1 \dots \bar{a}_{k-1} B} \eta^B$ and apply $L_{\bar{a}_k}$, using Lemma 14 and (21).

The proof of (26) is similar: We start with (18) for the case $k=1$, and for $k>1$, we use induction, Lemma 14, (20), and (21). $\square$

Proof of Theorem 18. Let $\mathbf{A} = (\mathbb{A}_1, \dots, \mathbb{A}_{\hat{n}})$. We rewrite the $\hat{n}$ equations obtained from (25) by replacing $\mathbb{A}$ with $\mathbb{A}_1, \dots, \mathbb{A}_{\hat{n}}$ in the following form:

$$(27) \quad M_{\mathbf{A}} \eta = w_{\mathbf{A}},$$

where

$$\eta = \begin{pmatrix} \eta^1 \\ \vdots \\ \eta^{\hat{n}} \end{pmatrix}, \quad w = \begin{pmatrix} w_{\mathbb{A}_1} \\ \vdots \\ w_{\mathbb{A}_{\hat{n}}} \end{pmatrix},$$

and the (i, B) th entry of $M_{\mathbf{A}}$ is

$$g_{\mathbb{A}_i B} - \sum_{\ell(\mathbb{A}') < \ell(\mathbb{A}_i)} g_{\mathbb{A}' B} v_{\mathbb{A}_i}^{\mathbb{A}'}.$$

Similarly, for each $a \in \{1, \dots, n\}$ we rewrite (26) as

$$(28) \quad N_{a, \mathbf{A}} \gamma_a = K_{a, \mathbf{A}} \eta + r_{\mathbf{A}}.$$

By Lemma 16 and Lemma 19 the matrices $M_{\mathbf{A}}$ and $N_{a, \mathbf{A}}$ are polynomial in $\overline{L^J \gamma_c^D}$, with coefficients smooth on $M \times \hat{M}$ and they only depend on M and $\hat{M}$. We define $\Delta_{0, \mathbf{A}} = \det M_{\mathbf{A}}$ and $\Delta_{j, \mathbf{A}} = \det N_{j, \mathbf{A}}$, $j = 1, \dots, n$, where we evaluate the coefficients of $\overline{L^J \gamma_c^D}$ at $(0, 0)$, and for $\mathbf{A} = (\mathbf{A}_0, \dots, \mathbf{A}_n)$, we set $\Delta_{\mathbf{A}} = \prod_{j=0}^n \Delta_{j, \mathbf{A}_j}$. By Cramer's rule, if $\Delta_{\mathbf{A}} \left(\overline{L^J \gamma_c^D}(0) \right) \neq 0$, in a neighbourhood of 0 we can solve the equations (27) with $\mathbf{A} = \mathbf{A}_0$ and (28) with $(a, \mathbf{A}) = (a_1, \mathbf{A}_1), \dots, (a_n, \mathbf{A}_n)$ for η^B and γ_a^B in the required form.

We finish the proof by showing that if f is k -nondegenerate at 0 with $k \leq k_0$, then there exists $\mathbf{A}^0$ with $\Delta_{\mathbf{A}^0} \left(\overline{L^J \gamma_c^D}(0) \right) \neq 0$. First recall that by Lemma 15, f is k -nondegenerate at 0 if and only if

$$(29) \quad \text{span}\{(g_{\mathbb{A}B}(0))_B : \ell(\mathbb{A}) \leq k\} = \mathbb{C}^{\hat{n}}.$$

Consider a system of vectors of the form

$$(30) \quad v_{\mathbb{A}} = \left(g_{\mathbb{A}B}(0) - \sum_{\ell(\mathbb{A}') < \ell(\mathbb{A})} c_{\mathbb{A}B}^{\mathbb{A}'} g_{\mathbb{A}'B}(0) \right)_B, \quad \ell(\mathbb{A}) \leq k.$$

We claim that if f is k -nondegenerate, then $\text{span}\{v_{\mathbb{A}} : \ell(\mathbb{A}) \leq k\} = \mathbb{C}^{\hat{n}}$, which implies that $\Delta_{\mathbf{A}^0} \left(\overline{L^J \gamma_c^D}(0) \right) \neq 0$ for some $\mathbf{A}^0$. To prove the last claim, we let $V_l = \text{span}\{(g_{\mathbb{A}B})_B(0) : \ell(\mathbb{A}) \leq l\}$ and $W_l = \text{span}\{v_{\mathbb{A}} : \ell(\mathbb{A}) \leq l\}$. By (30), $V_1 = W_1$. But then, again by (30), $V_2 = W_2$. By induction, we see that $W_k = V_k = \mathbb{C}^{\hat{n}}$. $\square$

Theorem 18 is completely analogous to Theorem 2.4 in [5]. The proof of Theorem 7 can now easily be completed using the arguments in [5] and a result from [8].

Proof of Theorem 7. By repeating the arguments in [5], based on the identities in Theorem 18 instead of those in Theorem 2.4 in [5], we obtain a complete system of some order l_0 (the exact value of which is not relevant for this argument) for f and g . That is, there exists a neighbourhood U of p_0 and a smooth function $\phi : U \times J^{l_0}(U, \hat{M}) \rightarrow J^{l_0+1}(U, \hat{M})$ such that

$$j_x^{l_0+1} f = \phi(x, j_x^{l_0} f) \text{ and } j_x^{l_0+1} g = \phi(x, j_x^{l_0} g), \quad x \in U.$$

By Theorem 2 in [8], $j_{p_0}^l f = j_{p_0}^l g$ for all l and, in particular then, $j_{p_0}^{l_0} f = j_{p_0}^{l_0} g$. The conclusion of Theorem 7 follows from the uniqueness of solutions of complete systems (see e.g. [4]). $\square$

5. JET REFLECTION IDENTITIES FOR LEVI-NONDEGENERATE HYPERSURFACES

In this section, we consider the case $\hat{N} = N + 1$, where M is Levi-nondegenerate at 0, and $\hat{M}$ is Levi-nondegenerate at $\hat{0}$. We are also assuming that f is transversal, which in our notation is equivalent to $\xi(0) \neq 0$.

For the proof of Theorem 1 we first note that by Lemma 20 in [8], the degeneracy of f is either 0 or 1. In the case where the degeneracy of f is 0, Theorem 1 follows from Theorem 7, so that we only have to deal with the case where f is constantly 1-degenerate. Using (18) it is easy to see that with our assumptions, the vectors $g_{aB}(0)$, $a = 1, \dots, n$ are linearly independent, as are the vectors $\gamma_c^D(0)$, $c = 1, \dots, n$. We conclude that f is constantly (1, 1)-degenerate. Our goal in this section is to derive the following jet reflection identity for such mappings. The unique determination theorem will be proved in section 7 by producing a complete system using this jet reflection identity.

Theorem 20. *Let $M \subset \mathbb{C}^N$ and $\hat{M} \subset \mathbb{C}^{N+1}$ be hypersurfaces which are Levi-nondegenerate at the points p_0 and $\hat{p}_0$, $f : M \rightarrow \hat{M}$ a smooth CR map which is constantly (1, 1)-degenerate and transversal at p_0 . Then for any I with $|I| = k$ there exist functions $r_b^{I,A}$ and $s^{I,A}$ for $a, b \in \{1, \dots, n\}$ and $A \in \{1, \dots, n+1\}$ such that*

$$(31) \quad L^I \gamma_b^A = r_b^{I,A} \left(L^M \gamma_c^D, L^M \eta^D, \overline{\gamma_c^D}, \overline{\eta^D} \right),$$

$$(32) \quad L^I \eta^A = s^{I,A} \left(L^M \gamma_c^D, L^M \eta^D, \overline{L_d \gamma_c^D}, \overline{L_d \eta^D}, \overline{T \gamma_c^D}, \overline{T \eta^D} \right)$$

Here $|M| \leq k - 1$. The $r_b^{I,A}$ and $s^{I,A}$ are rational in their arguments with coefficients which are smooth on $M \times \hat{M}$, and they only depend on M and $\hat{M}$.

Proof. Since $\hat{M}$ is Levi-nondegenerate, we can choose the basis of CR vector fields tangent to $\hat{M}$ such that

$$(33) \quad \hat{h}_{\bar{A}B}(0) = \epsilon_{\bar{A}B} = \begin{cases} \pm i, & A = B \\ 0, & A \neq B \end{cases}.$$

Since f is constantly $(1, 1)$ -degenerate, for any $\bar{a}, \bar{b}$ the determinant of the matrix

$$(34) \quad \begin{pmatrix} g_{\bar{1}1} & \cdots & g_{\bar{1}n+1} \\ \vdots & & \vdots \\ g_{\bar{n}1} & \cdots & g_{\bar{n}n+1} \\ L_{\bar{a}}g_{\bar{b}1} & \cdots & L_{\bar{a}}g_{\bar{b}n+1} \end{pmatrix}$$

vanishes. Let us write $(-1)^{B+1}\Delta^B$ for the determinant of the matrix obtained from the matrix (34) by dropping the last row and the B th column. We fix a and b for the moment. By developing the determinant of (34) along the last row,

$$(35) \quad \Delta^B L_{\bar{a}}g_{\bar{b}B} = 0.$$

We compute

$$(36) \quad L_{\bar{a}}g_{\bar{b}B} = L_{\bar{a}} \left(\overline{\gamma_b^A} \hat{h}_{\bar{A}B} \right) = \overline{L_a \gamma_b^A} \hat{h}_{\bar{A}B} + \overline{\gamma_b^A \gamma_a^D} \hat{L}_{\bar{D}} \hat{h}_{\bar{A}B}.$$

Plugging (36) into (35) and using Lemma 16 we conclude that

$$(37) \quad \Delta^B \overline{L_a \gamma_b^A} \hat{h}_{\bar{A}B} = r \left(\overline{\gamma_c^D} \right),$$

where r is rational in $\overline{\gamma_c^D}$ with coefficients smooth on $M \times \hat{M}$ and r is determined only by M and $\hat{M}$. We take the complex conjugate of this equation to obtain

$$(38) \quad \Delta^{\bar{B}} L_a \gamma_b^A \hat{h}_{A\bar{B}} = \bar{r} \left(\gamma_c^D \right),$$

Consider now the equations (18) which we rewrite as

$$\xi h_{\bar{d}b} = \gamma_b^A g_{\bar{d}A}.$$

We apply L_a to this equation and conclude, using also equations (18) and (19),

$$(39) \quad (L_a \gamma_b^A) g_{\bar{d}A} = \tilde{p}_{\bar{d}} \left(\gamma_c^D, \overline{\gamma_c^D}, \xi, L_a \xi \right) = p_{\bar{d}} \left(\gamma_c^D, \eta^D \overline{\gamma_c^D}, \overline{\eta^D} \right),$$

where $p_{\bar{d}}$ is rational in its arguments with coefficients smooth on $M \times \hat{M}$ and $p_{\bar{d}}$ is determined only by M and $\hat{M}$. We claim that the matrix

$$(40) \quad \begin{pmatrix} g_{\bar{1}1}(0) & \cdots & g_{\bar{1}n+1}(0) \\ \vdots & & \vdots \\ g_{\bar{n}1}(0) & \cdots & g_{\bar{n}n+1}(0) \\ \Delta^{\bar{1}}(0)\epsilon_{\bar{1}1} & \cdots & \Delta^{\overline{n+1}}(0)\epsilon_{\overline{n+1}n+1} \end{pmatrix}$$

is invertible, so that the system of equations

$$(41) \quad X^A g_{\bar{d}A} = p_{\bar{d}}, \quad d \in \{1, \dots, n\},$$

$$(42) \quad X^A \Delta^{\bar{B}} \hat{h}_{A\bar{B}} = \bar{r},$$

can be solved uniquely for the X^A (in a neighbourhood of 0). Developing the determinant Δ for the matrix (40) along the last row, we conclude that

$$(43) \quad \Delta = \epsilon_{\bar{A}A} \Delta^{\bar{A}} \Delta^A.$$

We are now going to show that this last expression is not equal to 0. Consider the matrices

$$(44) \quad G = \begin{pmatrix} g_{\bar{1}1}(0) & \cdots & g_{\bar{1}n+1}(0) \\ \vdots & & \vdots \\ g_{\bar{n}1}(0) & \cdots & g_{\bar{n}n+1}(0) \end{pmatrix}, \quad \Gamma = \begin{pmatrix} \gamma_{\bar{1}}^1(0) & \cdots & \gamma_{\bar{n}}^1(0) \\ \vdots & & \vdots \\ \gamma_{\bar{1}}^{n+1}(0) & \cdots & \gamma_{\bar{n}}^{n+1}(0) \end{pmatrix}$$

and the $n \times n$ matrix

$$(45) \quad H = \begin{pmatrix} h_{\bar{1}1}(0) & \cdots & h_{\bar{1}n}(0) \\ \vdots & & \vdots \\ h_{\bar{n}1}(0) & \cdots & h_{\bar{n}n}(0) \end{pmatrix}.$$

In matrix notation, (18) then reads as $\xi H = G\Gamma$. Let us write G_A for the matrix G with the A th column dropped, and Γ^A for the matrix Γ with the A th row dropped. The Cauchy-Binet Theorem from linear algebra states that $\det G\Gamma^t = \det G_A \det \Gamma^A$. By our convention (33) and (11) it follows that

$$g_{\bar{a}B}(0) = \overline{\gamma_a^B}(0) \epsilon_{BB}.$$

Writing $\epsilon = \prod_B \epsilon_{\bar{B}B}$ we obtain from the equation above

$$\Delta^A = \frac{\epsilon}{\epsilon_{\bar{A}A}} \det \overline{\Gamma^A}.$$

Recalling that $\Delta^A = \det G_A$, we see that the Cauchy-Binet Theorem implies that

$$\begin{aligned} \Delta &= \epsilon_{\bar{A}A} \Delta^{\bar{A}} \Delta^A \\ &= \epsilon \det \overline{G_A} \det \overline{\Gamma^A} \\ &= \epsilon \xi(0)^n \det H \neq 0, \end{aligned}$$

and our last claim is proved. Using that the matrix (40) is invertible to solve (38) and (39) for $L_a \gamma_b^A$, we obtain (by repeatedly differentiating the resulting equations and using the identities (20), (21), and (22)) the first part of Theorem 20.

We also need reflection identities (32) for $L_a \eta^B$. We will deduce those from the equation (31) we just proved. We first compute

$$L_{\bar{c}} L_b \gamma_e^B = L_b L_{\bar{c}} \gamma_e^B - h_{\bar{c}b} T \gamma_e^B,$$

which follows from (9) and the fact that $R_{\bar{a}c}^b = 0$ (cf. [5], Lemma 1.33). By (20) and (22), we conclude that

$$(46) \quad L_{\bar{c}} L_b \gamma_e^B = -L_b (\eta^B h_{\bar{c}e}) - h_{\bar{c}b} (L_e \eta^B + \eta^B h_e) = -(h_{\bar{c}e} L_b \eta^B + h_{\bar{c}b} L_e \eta^B) + r(\eta^B),$$

where $r(\eta^B)$ is a (linear) polynomial in η^B with coefficients which are smooth on M (and only depend on M). The identity (32) with $k = 1$ now follows by solving for $L_b \eta^B$ in (46) (e.g. set $b = c = e$ and recall that $h_{\bar{b}b} \neq 0$) and then applying $L_{\bar{c}}$ to (31) with $k = 1$, and finally using (20) and (21) to substitute for $L_{\bar{c}} \gamma_e^B$ and $L_{\bar{c}} \eta^B$. The general case follows by differentiating. The details are left to the reader. $\square$

6. STRICTLY PSEUDOCONVEX TARGETS

Our goal in this section is to derive the following reflection identity; in section 7 we will use it to derive a complete system for the mapping f .

Theorem 21. *Let $M \subset \mathbb{C}^N$, $\hat{M} \subset \mathbb{C}^{\hat{N}}$ be smooth hypersurfaces, $p_0 \in M$, $\hat{p}_0 \in \hat{M}$, with $\hat{M}$ strictly pseudoconvex at $\hat{p}_0$. Also assume that f is a smooth CR mapping from M to $\hat{M}$ with $f(p_0) = \hat{p}_0$ which is transversal and constantly (k_0, s) -degenerate at p_0 . Then for any I with $|I| = k$, $a \in \{1, \dots, n\}$, $B \in \{1, \dots, \hat{n}\}$ there exist functions $r_a^{I,B}$ and $s^{I,B}$ which are rational in their arguments such that*

$$(47) \quad L^I \gamma_a^B = r_a^{I,B} \left(L^M T^l \gamma_c^D, L^M T^l \eta^D, \overline{L^J T^m \gamma_c^D}, \overline{L^J T^m \eta^D} \right),$$

$$(48) \quad L^I \eta^B = s^{I,B} \left(L^M T^l \gamma_c^D, L^M T^l \eta^D, \overline{L^J T^m \gamma_c^D}, \overline{L^J T^m \eta^D} \right).$$

Here $l \in \{0, 1\}$, $|M| + l \leq \max(k_0, k-1)$, $|J| + m \leq k_0 + 1$, $m \leq \min(k, k_0)$. Furthermore, $r_a^{I,B}$ and $s^{I,B}$ only depend on M , $\hat{M}$, and $j_{p_0}^{k_0} f$.

Proof. Assuming that $\hat{M}$ is strictly pseudoconvex, we choose a local basis of CR vector fields tangent to $\hat{M}$ such that $\hat{h}_{\bar{A}B}(0) = i\delta_A^B$. Since f is transversal, it follows that M is strictly pseudoconvex at 0, and we likewise choose a local basis of CR vector tangent to M such that $h_{\bar{a}b}(0) = i\delta_a^b$. As in Theorem 2, we assume that f is constantly (k_0, s) -degenerate at p_0 . We let

$$V_k = \text{span}\{g_{\bar{A}B}(0) : \ell(\bar{A}) \leq k\}$$

and claim that we can assume that $V_{k_0} \subset \mathbb{C}^{\hat{n}}$ is just the subspace spanned by the first s unit vectors. To see this, recall that

$$g_{\bar{a}_1 \dots \bar{a}_k B} = \langle D_{\bar{a}_k} D_{\bar{a}_{k-1}} \dots D_{\bar{a}_1} \tau, \hat{L}_B \rangle.$$

Hence, under a change of basis $\hat{L}'_B = u_B^C \hat{L}_C$, the g transform as $g'_{\bar{A}B} = u_B^D g_{\bar{A}D}$. Also, since

$$\hat{h}_{\bar{A}B} = \langle d\hat{\theta}, \hat{L}_{\bar{A}} \wedge \hat{L}_B \rangle,$$

the $\hat{h}_{\bar{A}B}$ transform as $\hat{h}'_{\bar{A}B} = \overline{u_A^C} u_B^D \hat{h}_{\bar{C}D}$, and any u_A^B which is unitary at 0 will respect the normalization condition $\hat{h}_{\bar{A}B}(0) = i\delta_A^B$. So, our claim is proved.

Since f is constantly (k_0, s) -degenerate, we can choose $\mathbb{A}_{1,a}, \dots, \mathbb{A}_{s,a}$ for $a \in \{1, \dots, n\}$ and $\mathbb{A}_1, \dots, \mathbb{A}_s$, whose length does not exceed k_0 , with such that for any $a \in \{1, \dots, n\}$ the vectors

$$v_{a,j} = \left(g_{\bar{A}_j, aB} - \sum_{\ell(\bar{A}') < \ell(\bar{A}_j, a)} g_{\bar{A}'B} v_{\bar{A}_j}^{\bar{A}'} \left(\overline{L^J \gamma_c^D} \right) \right), \quad j = 1, \dots, s,$$

as well as

$$v_j = \left(g_{\bar{A}_j B} - \sum_{\ell(\bar{A}') < \ell(\bar{A}_j)} g_{\bar{A}'B} v_{\bar{A}_j}^{\bar{A}'} \left(\overline{L^J \gamma_c^D} \right) \right), \quad j = 1, \dots, s,$$

are linearly independent at 0, and hence each of these $n+1$ families of vectors span V_{k_0} . For any $t \in \{1, \dots, n\}$, $C \in \{1, \dots, \hat{n}\}$, and $I = (I_1, \dots, I_k)$, the determinants of

$$(49) \quad \begin{pmatrix} v_{a,1}^1 & \dots & v_{a,1}^s & v_{a,1}^C \\ \vdots & & \vdots & \vdots \\ v_{a,s}^1 & \dots & v_{a,s}^s & v_{a,s}^C \\ L^{\bar{I}} g_{\bar{a}1} & \dots & L^{\bar{I}} g_{\bar{a}s} & L^{\bar{I}} g_{\bar{a}C} \end{pmatrix}, \quad a = 1, \dots, n,$$

and

$$(50) \quad \begin{pmatrix} v_1^1 & \dots & v_1^s & v_1^C \\ \vdots & & \vdots & \vdots \\ v_s^1 & \dots & v_s^s & v_s^C \\ L^I g_{\bar{t}1} & \dots & L^I g_{\bar{t}s} & L^I g_{\bar{t}C} \end{pmatrix}$$

vanish on M . Since

$$L^I g_{\bar{t}B} = \overline{L^I (\gamma_t^A \hat{h}_{AB})} = \overline{L^I \gamma_t^A} \hat{h}_{AB} + q_{t,B}^I (\overline{L^J \gamma_c^D}),$$

where $|J| < k$, expanding (49) and (50) along the last row we obtain

$$(51) \quad \Delta_{\bar{a}, C_0}^{\bar{b}} (L^I \gamma_a^A) \hat{h}_{A\bar{b}} + \bar{\Delta}_a (L^I \gamma_a^A) \hat{h}_{A\bar{C}_0} = r_{a, C_0}^I (L^J \gamma_a^B), \quad a = 1, \dots, n,$$

$$(52) \quad \Delta_{\bar{C}_0}^{\bar{b}} (L^I \gamma_t^A) \hat{h}_{A\bar{b}} + \bar{\Delta} (L^I \gamma_t^A) \hat{h}_{A\bar{C}_0} = r_{C_0}^I (L^J \gamma_a^B),$$

where again $|J| < k$, r_{a, C_0}^I and $r_{C_0}^I$ are polynomial in their arguments with smooth coefficients, and $\Delta_{\bar{a}, C_0}^{\bar{b}}$, $\bar{\Delta}_a$, $\Delta_{\bar{C}_0}^{\bar{b}}$, $\bar{\Delta}$ denotes the various cofactors produced by the row expansion. These are rational in $L^J \gamma_a^B$, where $|J| \leq k_0 - 1$, and polynomial in $L^K \xi$, where $|K| \leq k_0$.

We need the following computational fact, which the reader can verify easily.

Lemma 22. *Given $P = p_1 \dots p_k$ with $1 \leq p_j \leq n$, the following holds for any a .*

$$(53) \quad L_{\bar{a}} L^P = L^P L_{\bar{a}} - \sum_{j=1}^k h_{\bar{a}p_j} L^{P_j} T + \sum_{|K| \leq k-2} C_K L^K T,$$

where P_j is obtained from P by removing the j th entry.

Now given any $I = i_1 \dots i_k$ with $i_1 \leq i_2 \leq \dots \leq i_k$, we first apply the operator L^I to the equations (25) and (26) to obtain

$$(54) \quad g_{\mathbb{A}B} (L^I \eta^B) - \sum_{\ell(\mathbb{A}') < \ell(\mathbb{A})} g_{\mathbb{A}'B} (L^I \eta^B) v_{\mathbb{A}}^{\mathbb{A}'} (\overline{L^J \gamma_c^D}) = \tilde{w}_{\mathbb{A}} (L^M \gamma_c^D, \overline{L^J T^m \gamma_c^D}, \overline{L^J T^m \eta^D})$$

and

$$(55) \quad g_{\mathbb{A}B} (L^I \gamma_a^B) - \sum_{\ell(\mathbb{A}') < \ell(\mathbb{A})} g_{\mathbb{A}'B} (L^I \gamma_a^B) p_{a\mathbb{A}}^{\mathbb{A}'} (\overline{L^J \gamma_c^D}) - \sum_{\ell(\mathbb{A}') < \ell(\mathbb{A})} g_{\mathbb{A}'B} (L^I \eta^B) q_{a\mathbb{A}}^{\mathbb{A}'} (\overline{L^J \gamma_c^D}) = \tilde{r}_{a\mathbb{A}} (L^M \gamma_c^D, \overline{L^J T^m \gamma_c^D}, \overline{L^J T^m \eta^D}),$$

where $|M| < k$, $|K| \leq k_0 - 1$, $|J| \leq k_0$, $m \leq k$, and $|J| + m \leq k_0$. We have also simplified matters a bit by applying (18) in order to get rid of the terms involving ξ .

Let us recall (52):

$$\Delta_{\bar{C}_0}^{\bar{b}} (L^I \gamma_t^A) \hat{h}_{A\bar{b}} + \bar{\Delta} (L^I \gamma_t^A) \hat{h}_{A\bar{C}_0} = r_{C_0}^I (L^M \gamma_a^B),$$

where $|M| \leq k-1$. Applying $L_{\bar{t}}$ to this equation, and applying Lemma 22 and (22), we obtain

$$\begin{aligned}
 (56) \quad & (L^I \gamma_t^A) L_{\bar{t}} \left(\Delta_{\bar{C}_0}^{\bar{b}} \hat{h}_{A\bar{b}} \right) + (L^I \gamma_t^A) L_{\bar{t}} \left(\bar{\Delta} \hat{h}_{A\bar{C}_0} \right) - \Delta_{\bar{C}_0}^{\bar{b}} (L^I \eta^A) h_{\bar{t}t} \hat{h}_{A\bar{b}} \\
 & - \bar{\Delta} (L^I \eta^A) h_{\bar{t}t} \hat{h}_{A\bar{C}_0} - \sum_{j=1}^k h_{\bar{t}i_j} (L^{I_j} L_t \eta^A) \Delta_{\bar{C}_0}^{\bar{b}} \hat{h}_{A\bar{b}} - \sum_{j=1}^k h_{\bar{t}i_j} (L^{I_j} L_t \eta^A) \bar{\Delta} \hat{h}_{A\bar{C}_0} \\
 & = \tilde{r}_{C_0}^I (L^M \eta^B, L^M \gamma_a^B, L^J T \eta^B, L^J T \gamma_a^B).
 \end{aligned}$$

In this equation, $|M| < k$, and $|J| < k-1$.

We consider the system of linear equations in the $L^I \gamma_a^B$ and $L^I \eta^B$ where I ranges over all $I = i_1 \dots i_k$ with $i_1 \leq i_2 \leq \dots \leq i_k$, $a \in \{1, \dots, n\}$, $B \in \{1, \dots, \hat{n}\}$ which is obtained in the following way. In (54) we set $\mathbb{A} = \mathbb{A}_1, \dots, \mathbb{A}_s$. For each I , this gives s linear equations in $L^I \eta^B$. We next consider the $\hat{n} - s$ equations obtained from (56) by replacing t by i_k and letting $C_0 = s+1, s+2, \dots, \hat{n}$. For the equations in $L^I \gamma_a^B$ ($a \in \{1, \dots, n\}$ fixed) we replace $\mathbb{A}$ in (55) by $\mathbb{A}_{a,1} \dots \mathbb{A}_{a,s}$, which gives us (for each I and each a) s linear equations in $L^I \gamma_a^B$. For each I and each a we also consider the equations obtained from (52) by replacing t by a , and letting $C_0 = s+1, \dots, \hat{n}$.

With the simplifications above, it is easy to see that this linear system can be solved for $L^I \gamma_a^B$ and $L^I \eta^B$. We can now lift the restriction that I is increasing by noting that rearranging the order of the i_j s only produces error terms of order less than k . This finishes the proof. $\square$

7. CONSTRUCTING THE COMPLETE SYSTEM; PROOF OF THEOREM 8

We are now going to construct the complete system for the jets of f . Since Theorems 20 and 21 are a bit different than the reflection identities in [5], we will go into some detail. For the rest of this section we assume that we are either in the situation of Theorem 20 or in the situation of Theorem 21. First, let us recall the following fact from [5] (the second part of Proposition 3.18):

For any multi-index J , integer $k \geq 1$, and index $c \in \{1, \dots, n\}$ there exist smooth functions $b_s^{e_1 \dots e_m}$ such that

$$(57) \quad \sum_{m=1}^{|J|+k} \sum_{s=0}^k b_s^{e_1 \dots e_m} \underbrace{[\dots [L_{e_1} \dots L_{e_m}, L_{\bar{c}}], L_{\bar{c}}] \dots, L_{\bar{c}}]}_{\text{length } s} = (h_{\bar{c}1})^p L^J T^k.$$

Here, $p = k + |J| - |J|_1 + 1$, $|J|_1$ denotes the number of occurrences of the index 1 in the multi-index J , and the length of the commutator $[\dots [X, Y_1], Y_2] \dots, Y_s]$ is s . Since in any case M is Levi-nondegenerate at 0, we can use this identity in order to find a formula for $L^J T^k \gamma_a^B$ and $L^J T^k \eta^B$. We first apply $L_{\bar{c}}$ (at most k times) to γ_a^B and η_a^B , and using equations (20) and (21) we see that the result is polynomial in γ_c^D , η_c^D , and $\overline{L_{\bar{c}}^j \gamma_c^D}$, where $j \leq k-1$. Next, we apply $L_{e_1} \dots L_{e_m}$ to the result, and we apply our reflection identities (either (31) and (32) or (47) and (48)) to see that the result is rational in

$$L^M T^l \gamma_c^D, L^M T^l \eta^D, \overline{L^K T^m \gamma_c^D}, \overline{L^K T^m \eta^D},$$

where $|M| + l \leq \max(k_0, |J| + k - 1)$, $|K| + m \leq k_0 + 1$, and $m \leq |J|$. Another application of $L_{\bar{c}}$ (at most k times) then gives us that

$$(58) \quad L^J T^k \gamma_a^B = u(L^M T^l \gamma_c^D, L^M T^l \eta^D, \overline{L^K T^m \gamma_c^D}, \overline{L^K T^m \eta^D}),$$

$$(59) \quad L^J T^k \eta^B = v(L^M T^l \gamma_c^D, L^M T^l \eta^D, \overline{L^K T^m \gamma_c^D}, \overline{L^K T^m \eta^D}),$$

where $|M| + l \leq \max(k_0, |J| + k - 1)$, $|K| + m \leq k + k_0 + 1$, and $m \leq \min(|J| + k, k_0)$. We are abusing notation by denoting by u , v and w functions which are rational in their arguments with coefficients smooth on $M \times \hat{M}$ and which only depend on M and $\hat{M}$ (and are allowed to change). Now in these

equations for $|J| + k = 2k_0 + 2$ we can substitute for the conjugated terms, yielding the same equations, but with the bounds $|M| + l \leq 2k_0 + 1$, $|K| + m \leq 2k_0 + 1$. Another use of equations (20) and (21) lets us conclude that for all J, k, R with $|J| + k + |R| = 2k_0 + 2$,

$$(60) \quad L^J T^k L^{\bar{R}} \gamma_a^B = u(L^M T^l \gamma_c^D, L^M T^l \eta^D, \overline{L^K T^m \gamma_c^D}, \overline{L^K T^m \eta^D}),$$

$$(61) \quad L^J T^k L^{\bar{R}} \eta^B = v(L^M T^l \gamma_c^D, L^M T^l \eta^D, \overline{L^K T^m \gamma_c^D}, \overline{L^K T^m \eta^D}),$$

with $|M| + l \leq 2k_0 + 1$, $|K| + m \leq 2k_0 + 1$. These two equations together with (18) let us also see that for all J, k, R with $|J| + k + |R| = 2k_0 + 2$,

$$(62) \quad L^J T^k L^{\bar{R}} \xi = w(L^M T^l \gamma_c^D, L^M T^l \eta^D, \overline{L^K T^m \gamma_c^D}, \overline{L^K T^m \eta^D}),$$

again with $|M| + l \leq 2k_0 + 1$, $|K| + m \leq 2k_0 + 1$. Equations (60), (61), and (62), written in some local coordinate systems, constitute the complete system of differential equations described in Theorem 8. Also note that the same complete system works for mappings whose jets are close to the $2k_0 + 2$ jet of f . This finishes the proof of Theorem 8.

REFERENCES

- [1] M. S. Baouendi, P. Ebenfelt, and L. P. Rothschild. Local geometric properties of real submanifolds in complex space. *Bull. Amer. Math. Soc. (N.S.)*, 37(3):309–336 (electronic), 2000.
- [2] M. S. Baouendi, P. Ebenfelt, and L. P. Rothschild. *Real submanifolds in complex space and their mappings*. Princeton University Press, Princeton, NJ, 1999.
- [3] M. S. Baouendi, X. Huang, and L. P. Rothschild. Regularity of CR mappings between algebraic hypersurfaces. *Invent. Math.*, 125(1):13–36, 1996.
- [4] R. L. Bryant, S. S. Chern, R. B. Gardner, H. L. Goldschmidt, and P. A. Griffiths. *Exterior Differential Systems*, volume 18 of *Mathematical Sciences Research Institute Publications*. Springer-Verlag, New York, 1991.
- [5] P. Ebenfelt. Finite jet determination of biholomorphic mappings at the boundary. To appear, *Asian J. Math.*
- [6] F. Forstnerič. Extending proper holomorphic mappings of positive codimension. *Invent. Math.*, 95(1):31–61, 1989.
- [7] P. Griffiths and J. Harris. *Principles of algebraic geometry*. John Wiley & Sons Inc., New York, 1994. Reprint of the 1978 original.
- [8] B. Lamel. Holomorphic maps of real submanifolds in complex spaces of different dimensions. *Pacific J. Math.*, 201(2):357–387, 2001.
- [9] B. Lamel. A reflection principle for real-analytic submanifolds of complex spaces. *J. Geom. Anal.*, 11(4):625–631, 2001.
- [10] D. Zaitsev and S.-Y. Kim. The equivalence and the embedding problems for cr-structures of any codimension. Preprint, math.CV/0108093, 2001.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT SAN DIEGO, LA JOLLA, CA 92093
E-mail address: `pebenfel@math.ucsd.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA
E-mail address: `lamelb@member.ams.org`